

1. Fix $x > 0$. Prove $(1 + x)^n \geq 1 + nx$ for any $n \in \mathbb{N}$. Deduce that $(1 + x)^{-n} \rightarrow 0$. Deduce that if $r \in (0, 1)$ then $r^n \rightarrow 0$, and if $r \in (1, \infty)$ then $r^n \rightarrow \infty$.
2. Suppose $\lim_{n \rightarrow \infty} |a_{n+1}/a_n| = L$. In lectures we proved that if $L < 1$ then $a_n \rightarrow 0$.
 - (a) Prove that if $L > 1$ then $|a_n| \rightarrow \infty$.
 - (b) Give an example with $|a_{n+1}/a_n| < 1 \forall n$ but $a_n \not\rightarrow 0$.

Give (without proof) examples where $L = 1$ and

- | | |
|-----------------------------------|------------------------------------|
| (i) $a_n \rightarrow 0$, | (iii) a_n divergent and bounded, |
| (ii) $a_n \rightarrow a \neq 0$, | (iv) $a_n \rightarrow \infty$. |

3. Let $(a_n)_{n \geq 1}$ be a sequence of *strictly positive* real numbers. Give an example such that $(1/a_n)_{n \geq 1}$ is unbounded. Suppose that $a_n \rightarrow a \neq 0$. Prove *from first principles* that $(1/a_n)_{n \geq 1}$ is bounded.
- 4.† Fix $r \in (0, 1/8)$. Define $(a_n)_{n \geq 1}$ by $a_1 := 1$ and $a_{n+1} = ra_n^2 + 1$.

- (a) Show that $a_{n+1} - a_n = r(a_n + a_{n-1})(a_n - a_{n-1})$.
- (b) Show that if $0 < a_j < 2 \quad \forall j \leq n$, (1)
then $|a_{n+1} - a_n| < (4r)^n/4$. (2)
- (c) Deduce that if (1) holds, then $a_{n+1} < r/(1 - 4r) + 1$.
- (d) Conclude that (1) holds for $j = n + 1$ too, and so $\forall j$ by induction.
- (e) Using (2) deduce $|a_m - a_n| < (4r)^n/4(1 - 4r)$ for $m \geq n$.
- (f) Deduce a_n is Cauchy. What does it converge to?

- 5.* Show that *any* sequence of real numbers $(a_n)_{n \geq 0}$ has a subsequence which either converges, or tends to ∞ , or tends to $-\infty$.

6. At home Professor Papageorgiou has made a fully realistic mathematical model of a dart board. It is a copy of the unit interval $[0, 1]$ in a frictionless vacuum. He throws a countably infinite number of darts at it, the n th landing at $a_n \in [0, 1]$.

He then makes a small dot $(x - \epsilon_x, x + \epsilon_x)$ around each point $x \in [0, 1]$ with his pen. Prove that however small he makes each dot, at least one of them will contain an infinite number of darts $a_n \in [0, 1]$.

What if he only makes dots around each dart $a_n \in [0, 1]$?

7. Let $(a_n)_{n \geq 1}$ be the sequence $\frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \frac{1}{6}, \dots$
 - (i) Give (without proof) a subsequence of $(a_n)_{n \geq 1}$ which converges to $\ell = 0$, and one which converges to $\ell = 1$.
 - (ii) Given any $\ell \in (0, 1)$, give (with proof) a subsequence convergent to ℓ .

8. A student is learning about Cauchy sequences, and thinks they have a brilliant proof that allows them to precisely identify the limit of a Cauchy sequence straight from the Cauchy condition. The student gives their proof below, is it correct?

$$\begin{aligned}\forall \epsilon > 0 \exists N \in \mathbb{N} \text{ such that } n, m \geq N &\Rightarrow |a_n - a_m| < \epsilon \\ \Rightarrow \forall n \geq N \quad |a_n - a_N| < \epsilon & \\ \Rightarrow a_n \rightarrow a_N \text{ as } n \rightarrow \infty. &\end{aligned}$$

*Starred questions * are good to prepare to discuss at your Problem Class.*

Questions marked † are slightly harder (closer to exam standard), but good for you.