

1. Evaluate $\int_0^x \frac{1}{1+e^t} dt$. Does $\int_0^\infty \frac{1}{1+e^t} dt$ exist, and if so, what is it?
2. The prime number theorem says that the number $\pi(n)$ of primes between 1 and n is approximately $\int_2^n \frac{1}{\log(x)} dx$.
 - (a) Prove that this integral equals $\frac{n}{\log(n)} + \int_2^n \frac{1}{(\log x)^2} dx$, up to a constant which does not depend on n .
 - (b) Prove that there is a constant $C > 0$ such that $\int_2^n \frac{1}{(\log x)^2} dx < \frac{Cn}{(\log n)^2}$ for all sufficiently large n , by splitting the integral up into one with domain $[2, \sqrt{n}]$ and one with domain $[\sqrt{n}, n]$ and estimating each one separately.
3. Let $f : [0, \infty) \rightarrow [0, \infty)$ be uniformly continuous, and suppose that $\int_0^\infty f(x) dx$ exists.
 - (a) For each $\epsilon > 0$, prove that there is a $\delta > 0$ such that for all $y > 0$, if $f(y) \geq \epsilon$ then


$$\int_y^{y+\delta} f(t) dt \geq \frac{\epsilon\delta}{2}.$$
 - (b) Prove that $\lim_{x \rightarrow \infty} f(x) = 0$.
 - (c) Describe a continuous function $g : [0, \infty) \rightarrow [0, \infty)$ such that $\int_0^\infty g(x) dx$ exists but $\lim_{x \rightarrow \infty} g(x)$ does not. Can you make g differentiable as well?
4. Let $\Gamma(t) = \int_0^\infty x^{t-1} e^{-x} dx$.
 - (a) Prove that this improper integral converges for all $t > 0$. (In how many ways is it improper?)
 - (b) Compute $\Gamma(1)$.
 - (c) Prove that $\Gamma(n+1) = n\Gamma(n)$ for all integers $n \geq 1$, and deduce that $\Gamma(n+1) = n!$ for all $n \geq 0$.
5. (*) Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous, with $f''(x)$ continuous and bounded on (a, b) .
 - (a) Use integration by parts twice to prove that

$$\int_a^b \frac{(x-a)(x-b)}{2} f''(x) dx = \int_a^b f(x) dx - (b-a) \left(\frac{f(a) + f(b)}{2} \right).$$

- (b) If $|f''(x)| \leq M$ for all $x \in (a, b)$, prove that

$$\left| \int_a^b \frac{(x-a)(x-b)}{2} f''(x) dx \right| \leq \frac{M(b-a)^3}{12}.$$

In other words, $\int_a^b f(x) dx$ is the area of the trapezium shown at right, up to an error of at most $\frac{M(b-a)^3}{12}$. (Hint: check that $(x-a)(x-b) \leq 0$ on $[a, b]$, and compute that $\int_a^b (x-a)(x-b) dx = -\frac{(b-a)^3}{6}$.)



(c) Apply this to $f(x) = \log(x)$ to show that

$$\int_1^n \log(x) dx = \sum_{k=1}^{n-1} \left(\frac{\log(k) + \log(k+1)}{2} + e_k \right),$$

where $|e_k| \leq \frac{1}{12k^2}$ for all k .

(d) Evaluate both the integral and the sum from part (c) to show that there is some constant $C > 0$ such that

$$\left| \log(n!) - \log \left(\frac{n^{n+1/2}}{e^{n-1}} \right) \right| < C$$

for all n , or equivalently if $C_1 = e^{1-C}$ and $C_2 = e^{1+C}$ then

$$C_1 \sqrt{n} \left(\frac{n}{e} \right)^n \leq n! < C_2 \sqrt{n} \left(\frac{n}{e} \right)^n.$$