

# Mathematics Year 1, Calculus and Applications I

D.T. Papageorgiou

## Problem Sheet 4

- (a) Show that  $\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$ . [Hint: Consider  $\sum_{i=1}^n [(i+1)^3 - i^3]$ ]  
(b) Find the integral  $\int_0^1 x^2 dx$  using upper Riemann sums and an equipartition of  $[0, 1]$ .
- In approximating the integral  $\int_0^1 e^x dx$  with an upper Riemann sum, we used the result  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} e^{i/n} = e - 1$ . Show this.

- Show that the function

$$f(x) = \begin{cases} 0 & \text{if } x \text{ is rational} \\ 1 & \text{if } x \text{ is irrational} \end{cases} ,$$

is not Riemann integrable. [Hint: Take *any* partition of  $[0, 1]$  and consider the lower and upper Riemann sums.]

- Show that  $\frac{d}{dx}(\sec x + \tan x) = \sec x(\sec x + \tan x)$ . Hence show that

$$\int \sec x dx = \log(\sec x + \tan x).$$

Note that the integral derived above makes sense only if (i)  $\cos x \neq 0$ , and (ii)  $\sec x + \tan x > 0$ . Determine an interval where such an interval can be applied.

- Calculate

$$\int \frac{1}{(x^2+1)^3} dx, \quad \int \frac{1}{x^3-1} dx, \quad \int \frac{x^3+1}{x^3-1} dx, \quad \int x^3 \sqrt{x^2+1} dx, \quad \int_{\pi/6}^{\pi/2} \frac{\cos x}{\sin x + \sin^3 x} dx$$

- Let  $I_n = \int \frac{1}{(x^2+1)^n} dx$  where  $n > 1$  is an integer (what is the integral when  $n = 1$ ?) Starting from  $I_{n-1}$  use integration by parts to establish the recursion formula

$$2(n-1)I_n = \frac{x}{(x^2+1)^{n-1}} + (2n-3)I_{n-1}.$$

- (a) Show that for any two integers  $m, n$

$$\int_{-\pi}^{\pi} \sin mx \cos nx dx = 0, \quad \int_{-\pi}^{\pi} \sin mx \sin nx dx = \begin{cases} 0 & \text{if } m \neq n \\ \pi & \text{if } n = m \end{cases} ,$$

and find an analogous formula for  $\int_{-\pi}^{\pi} \cos mx \cos nx dx$ .

- (b) Suppose that  $f$  is defined on  $[-\pi, \pi]$  and it is a  $2\pi$ -periodic function, i.e.  $f(x+2\pi) = f(x)$ . If we approximate  $f(x)$  by the series  $f(x) \approx a_0 + \sum_{k=1}^N a_k \cos kx + b_k \sin kx$ , use the results above to find formulas for  $a_0, a_k, b_k, k = 1, \dots, N$ .

(c) Now take  $f$  to be defined on  $[-\pi, \pi]$  as follows

$$f(x) = \begin{cases} 1 & \text{if } |x| \leq \pi/2 \\ 0 & \text{otherwise} \end{cases} .$$

Using the trigonometric approximation of part (b) above, calculate the a's and b's and confirm that  $b_k = 0$  for all  $k$ . Could you have anticipated this result by considering the symmetries of  $f$ ?

*Congratulations! You have just computed a Fourier series.*

8. Using comparison tests for improper integrals, determine convergence or divergence of the following integrals

$$\int_0^\infty e^{-x^2} dx, \quad \int_0^\infty \frac{x^3}{(1+x^2)^2} dx, \quad \int_0^\infty \frac{1}{\sqrt{x+x^3}} dx$$

$$\int_0^1 \frac{\sin^2 x}{1+x^2} dx, \quad \int_0^1 \frac{1}{\log(1+x)} dx, \quad \int_0^\infty \sin(x^2) dx.$$

9. Prove that

$$\int_0^1 \frac{x^3}{2 - \sin^4 x} dx \leq \frac{1}{4} \log 2 \quad \text{and} \quad \left| \int_0^{\pi/2} \frac{x - \pi/2}{2 + \cos x} dx \right| \leq \frac{\pi^2}{16}.$$

10. Prove the *integral mean value theorem* which generalizes a bit the theorem proved in class: Let  $f$  and  $g$  be continuous on  $[a, b]$  with  $g(x) \geq 0$  for  $x \in [a, b]$ . Then there exists a  $c$  between  $a$  and  $b$  with

$$\int_a^b f(x)g(x)dx = f(c) \int_a^b g(x)dx.$$

Show by finding an example that the conclusion of the theorem is wrong if the assumption  $g(x) \geq 0$  is dropped.

11. Let  $\mu$  be the average of the function  $f$  defined on a closed interval  $[a, b]$ . The average value of  $(f(x) - \mu)^2$  is called the *variance* of  $f$  on  $[a, b]$ , and the square root of the variance is the *standard deviation* of  $f$  on  $[a, b]$  and is denoted by  $\sigma$ . Find the average value, variance and standard deviation of each of the following functions on the specified interval:

$$x^2 \text{ on } [0, 1], \quad xe^x \text{ on } [0, 1], \quad \sin 2x \text{ on } [0, 4\pi],$$

$$f(x) = \begin{cases} 1 & \text{on } [0, 1] \\ 2 & \text{on } (1, 2] \end{cases}, \quad f(x) = \begin{cases} 2 & \text{on } [0, 1] \\ 3 & \text{on } (1, 2] \\ 1 & \text{on } (2, 3] \\ 5 & \text{on } (3, 4] \end{cases}$$

12. (a) Suppose that  $f(x)$  is a step function on  $[a, b]$ , with value  $k_i$  on the interval  $(x_{i-1}, x_i)$  belonging to the partition  $(x_0, x_1, \dots, x_n)$ . Find a formula for the standard deviation of  $f$  on  $[a, b]$ .
- (b) Simplify your formula for the case when the partition consists of equally spaced points.

- (c) Show that if the standard deviation of a step function is zero, then the function is a constant.
- (d) Give a definition of the standard deviation of a list of numbers  $a_1, a_2, \dots, a_n$ .
- (e) What can you say about the list of numbers if its standard deviation is zero?
13. Given a positive integer  $n$  define

$$\delta_n(x) = \begin{cases} n & \text{for } |x| \leq \frac{1}{2n} \\ 0 & \text{otherwise} \end{cases}$$

Consider also a continuous function  $g(x)$  defined on  $[a, b]$  where  $a < 0, b > 0$ .

- (a) Calculate  $\int_a^b g(x)\delta_n(x)dx$ , and prove that  $\lim_{n \rightarrow \infty} \int_a^b g(x)\delta_n(x)dx = g(0)$ .
- (b) Find also  $\lim_{n \rightarrow \infty} \int_a^t g(x)\delta_n(x)dx$  for  $t < 0$  and  $t > 0$ , respectively.
- (c) Given arbitrary numbers  $t_1, t_2$ , define  $\int_{t_1}^{t_2} \delta(x)dx$  by the  $\lim_{n \rightarrow \infty} \int_{t_1}^{t_2} \delta_n(x)dx$ . Use the results of part (b) above to find the former integral for  $t_1 < t_2 < 0, 0 < t_1 < t_2$  and  $t_1 < 0 < t_2$ .
- (d) The function  $\delta(x)$  is a *distribution* - it is zero everywhere except at 0 where it is infinite. What is its anti-derivative? Give an expression and sketch it.
14. Let  $f(x) = [x] + 1$  and  $F(x) = \int_0^x f(t)dt$  (recall that  $[x]$  means the integer part of  $x$ ). Find an explicit expression for  $F(x)$  when  $0 \leq x \leq 2$ , and show that  $F'(1) \neq f(1)$ . Explain why this does not contradict the fundamental theorem of calculus.
15. By evaluating the integral  $\int_1^n \log x dx$  where  $n$  is a positive integer, and comparing with the upper and lower Riemann sums associated to the partition  $(1, 2, \dots, n)$  of the interval  $[1, n]$ , show that

$$(n-1)! \leq n^n e^{-n} e \leq n!$$

Hence prove that

$$\lim_{n \rightarrow \infty} \left( \frac{n!}{n^n} \right)^{1/n} = 1/e.$$

16. For any non-negative integer  $n$ , let

$$I_n = \int_0^\infty e^{-x} (\sin x)^n dx, \quad J_n = \int_0^\infty e^{-x} (\sin x)^n \cos x dx.$$

- (a) Show that  $I_0 = 1, J_0 = 1/2, I_n = nJ_{n-1}$  and  $J_n = nI_{n-1} - (n+1)I_{n+1}$ .
- (b) Using the results in part (a), show that  $I_1 = 1/2, J_1 = 1/5$  and that for  $n \geq 2$  we have the explicit recursion formulas

$$I_n = \frac{n(n-1)}{(1+n^2)} I_{n-2}, \quad J_n = \frac{n(n-1)}{1+(n+1)^2} J_{n-2}, \quad n \geq 2.$$

- (c) Find explicit expressions in the form of rational numbers for each of  $I_n$  and  $J_n$ . [Note: You need to treat  $n$  being even or odd sep separately.] Which is larger,  $I_n$  or  $J_n$ ? Explain.