

MATH50001 Problems Sheet 4

Solutions

1)

$$\oint \frac{z^2}{(z-1)^n} dz = \frac{2\pi i}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} z^2 = \begin{cases} 2\pi i, & n=1, \\ 4\pi i, & n=2, \\ 2\pi i, & n=3, \\ 0, & n>3. \end{cases} .$$

2 a) This is the ellipse with two foci at 2 and -2 .

2 b)

$$\oint_{\gamma} \frac{\sin z}{(z+2)^3} dz = \frac{2\pi i}{2!} \frac{d^2}{dz^2} \sin z \Big|_{z=-2} = -\pi i \sin(-2) = \pi i \sin 2.$$

3) Let p be a polynomial. Then, by Cauchy's formula

$$\frac{1}{2\pi i} \oint_{|z|=1} \frac{1-zp(z)}{z} dz = \frac{1}{2\pi i} \oint_{|z|=1} \frac{1}{z} dz = 1.$$

Therefore by using the ML-inequality we obtain

$$1 = \left| \frac{1}{2\pi i} \oint_{|z|=1} \frac{1-zp(z)}{z} dz \right| \leq \max_{|z|=1} |1-zp(z)| = \max_{|z|=1} |z^{-1} - p(z)|.$$

4) Indeed, for any $z_0, z_1 \in \mathbb{C}$, $z_0 \neq z_1$ and R sufficiently large, we have

$$\begin{aligned} & \frac{1}{2\pi i} \oint_{|z|=R} \frac{f(z)}{(z-z_0)(z-z_1)} dz \\ &= \frac{1}{z_0-z_1} \left(\frac{1}{2\pi i} \oint_{|z|=R} \frac{f(z)}{(z-z_0)} dz - \frac{1}{2\pi i} \oint_{|z|=R} \frac{f(z)}{(z-z_1)} dz \right) \\ &= \frac{1}{z_0-z_1} (f(z_0) - f(z_1)). \end{aligned}$$

(3 unseen)

2

Since f is bounded, there is a constant M , such that $|f(z)| \leq M$, $z \in \mathbb{C}$. Therefore using the ML-inequality we find

$$\begin{aligned} \left| \frac{1}{2\pi i} \oint_{|z|=R} \frac{f(z)}{(z-z_0)(z-z_1)} dz \right| &\leq M R \max_{z:|z|=R} \frac{1}{|(z-z_0)(z-z_1)|} \\ &\leq M R \frac{1}{(R-|z_0|)(R-|z_1|)} \\ &= M R^{-1} \frac{1}{(1-|z_0|/R)(1-|z_1|/R)} \rightarrow 0, \end{aligned}$$

as $R \rightarrow \infty$

This implies

$$\frac{1}{z_1 - z_0} (f(z_0) - f(z_1)) = 0$$

and thus $f(z_0) = f(z_1)$. Since z_0 and z_1 are arbitrary, we finally obtain that f is a constant function.

5) Note that if $n = 0$, then we simply apply Liouville's theorem.

Assume that $|f(z)| \leq C(1 + |z|)^n$ with some $C > 0$. Then for any $z_0 \in \mathbb{C}$ we have

$$\begin{aligned} |f^{(n+1)}(z_0)| &= \left| \frac{(n+1)!}{2i\pi} \oint_{|z-z_0|=R} \frac{f(z)}{(z-z_0)^{n+2}} dz \right| \\ &\leq \frac{C(n+1)!}{2\pi} \max_{z:|z-z_0|=R} (1+|z|)^n \frac{2\pi R}{R^{n+2}} \rightarrow 0, \end{aligned}$$

as $R \rightarrow \infty$.

Therefore $f^{(n+1)} \equiv 0$ and thus $f^{(n)}$ is a constant function. We conclude that $f(z)$ is a polynomial of degree at most n by integrating $f^{(n)}(z)$ n -times.

6) Assume that $f = u + iv$ is an entire function that has a bounded real part. Then $g(z) = e^{f(z)}$ is also entire. Note that since u is bounded then $|g| = e^u$ is bounded. Thus g is constant. and therefore f is a constant function.

7)

- a) converges,
- b) converges,

$$\left| \frac{3 - (2i)^n}{\cos ni} \right| = 2 \left| \frac{3 - (2i)^n}{e^{-n} + e^n} \right| = 2 \frac{2^n |3/2^n - i^n|}{e^n (1 + e^{-2n})}$$

Clearly $|3/2^n - i^n| \leq 3/2^n + 1 \leq 5/2$ and $1 + e^{-2n} > 1$. Therefore

$$\left| \frac{3 - (2i)^n}{\cos ni} \right| \leq 5 \frac{2^n}{e^n}.$$

Since $2 < e$ we conclude that series converges.

c) diverges, indeed:

$$\left| \frac{ni}{n+i} \right|^{n^2} = \left(\frac{n}{\sqrt{n^2+1}} \right)^{n^2} = \left(\frac{1}{\sqrt{1+1/n^2}} \right)^{n^2} = \frac{1}{(1+1/n^2)^{n^2/2}} \rightarrow e^{-1/2} \neq 0.$$

Because it is known that

$$\lim_{t \rightarrow \infty} \left(1 + \frac{1}{t} \right)^t = e.$$

8) $\operatorname{Re} z \leq 0$.

9) a) $|z| < 1$; b) $|z - 4| < 2^{-1/4}$; c) $|z - 2| < 1$.

10)

a) $\sum_{n=2}^{\infty} \frac{z^n}{2^{n-1}}, \quad |z| < 2$

b) $\sum_{n=0}^{\infty} \frac{(-1)^n}{(1+i)^{n+1}} (z-i)^n, \quad |z-i| < \sqrt{2}$.

11)

a) $\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} z^{2n}, \quad |z| < \infty$.

b) $\sum_{n=0}^{\infty} \frac{e^{1+i}}{n!} (z-1-i)^n, \quad |z-1-i| < \infty$.

c) $\frac{\pi i}{2} + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{ni^n} (z-i)^n, \quad |z-i| < 1$.

12)

Clearly

$$a_k = \frac{1}{k!} f^{(k)}(0).$$

a) Let $1 > \varepsilon > 0$. By using the generalized Cauchy's formula we find

$$f^{(k)}(0) = \frac{k!}{2i\pi} \oint_{|z|=1-\varepsilon} \frac{f(z)}{z^{k+1}} dz.$$

Therefore by using the ML inequality and the fact that $|f(z)| < 1$ in \mathbb{D} we have

$$|a_k| \leq \frac{1}{k!} \left| \frac{k!}{2i\pi} \oint_{|z|=1-\varepsilon} \frac{f(z)}{z^{k+1}} dz \right| \leq \frac{1}{(1-\varepsilon)^k}.$$

Letting $\varepsilon \rightarrow 0$ we obtain $|a_k| \leq 1$.

b) We now use the generalized Cauchy's formula integrating over a circle $C_r = \{z \in \mathbb{C} : |z| = r\}$.

$$f^{(k)}(0) = \frac{k!}{2i\pi} \oint_{|z|=r} \frac{f(z)}{z^{k+1}} dz.$$

4

By applying the ML inequality we find by using the inequality $|f(z)| < (1 - |z|)^{-1}$

$$|a_k| \leq \frac{r}{r^{k+1}(1-r)}$$

Note that

$$\frac{d}{dr} r^k(1-r) = kr^{k-1} - (k+1)r^k = 0$$

implies $r = k/(k+1)$ and finally we obtain

$$|a_k| \leq \frac{(k+1)^{k+1}}{k^k}.$$